



# Transcriptomics exposes the uniqueness of parasitic plants

Yasunori Ichihashi, J. Musembi Mutuku, Satoko Yoshida and Ken Shirasu

Corresponding author. K. Shirasu, RIKEN Center for Sustainable Resource Science, 1-7-22 Suehiro-cho, Tsurumi-ku, Yokohama, Kanagawa 230-0045, Japan. Tel.: +81-45-503-9574; Fax: +81-45-503-9573. E-mail: ken.shirasu@riken.jp

## Abstract

Parasitic plants have the ability to obtain nutrients directly from other plants, and several species are serious biological threats to agriculture by parasitizing crops of high economic importance. The uniqueness of parasitic plants is characterized by the presence of a multicellular organ called a haustorium, which facilitates plant–plant interactions, and shutting down or reducing their own photosynthesis. Current technical advances in next-generation sequencing and bioinformatics have allowed us to dissect the molecular mechanisms behind the uniqueness of parasitic plants at the genome-wide level. In this review, we summarize recent key findings mainly in transcriptomics that will give us insights into the future direction of parasitic plant research.

**Key words:** parasitic plants; transcriptomics; haustorium; plant–plant interaction; photosynthesis

## Introduction

Parasitic plants acquire water, carbon and nutrients via vascular connections to the host plants [1, 2]. Parasitic plants consist of ~4000 species from 19 different families that can be found in most major biomes [3]. They can be classified based on their life cycle and mode of nutrition: (i) dependence on host: a parasite that requires a host to complete its life cycle is termed as 'obligate', a parasite that does not is termed as 'facultative'; (ii) presence or absence of chlorophyll: partially photosynthetic are termed as 'hemiparasitic', and non-photosynthetic as 'holoparasitic'; (iii) points of attachment: root or stem parasites [4]. Parasitism in angiosperms has originated independently at least 11 times [5], which is an example of convergent evolution.

Parasitic plants develop a multicellular organ called a haustorium, whose functions include attachment and invasion to a host and the physiological redirection of host resources into the parasite [2]. Haustorium formation and seed germination occur in response to host-derived chemical cues [2, 6]. Parasites and host plants exchange mobile molecules through the haustorium [7] and exhibit defense response at the haustorial interface in the root tissues, such as root endodermis, cortex and epidermal layers, which are in contact with the host [8, 9]. In extreme cases of plant parasitism (holoparasite), the parasites have lost their chloroplasts and are unable to carry out photosynthesis [10]. Thus, the uniqueness of parasitic plants is characterized by the development of the haustorium that enables nutrient acquisition and reduction of photosynthesis, contrasting with

**Yasunori Ichihashi** is a Special Postdoctoral Researcher at RIKEN Center for Sustainable Resource Science, Plant Immunity Research Group. He specializes in evolutionary developmental biology of plant shape and transcriptomics. Currently, he is working in genomics of parasitic plants.

**Josiah Musembi Mutuku** is a Japan Society for the Promotion of Science (JSPS) Postdoctoral Researcher at RIKEN Center for Sustainable Resource Science, Plant Immunity Research Group. He specializes in using molecular biology and genetics tools to characterize host resistance response to pathogens including fungi. Currently, he is studying defense signaling during host response to the root hemiparasite *Striga hermonthica*.

**Satoko Yoshida** is a Senior Research Scientist at RIKEN Center for Sustainable Resource Science, Plant Immunity Research Group. She specializes in molecular biology of plant parasitism and symbiosis. She currently focuses on genomics and genetics of parasitic plants, especially species in Orobanchaceae, such as *Striga* and *Phtheirospermum*.

**Ken Shirasu** is a Group Director at RIKEN Center for Sustainable Resource Science, Plant Immunity Research Group. He has been studying molecular plant biotic interactions. His current focus is on molecular mechanisms of plant immunity and pathogen factors that overcome the immunity system.

autotrophy of typical plants that produce organic nutrients through photosynthesis [2].

Several species of parasitic plants have severe effects on host plant growth and have a serious economic impact on agriculture [2, 4, 11]. Since agriculture began ~10 000 years ago, the invasion of parasitic plants from tropical grasslands to agricultural lands has occurred within a relatively short period [12]. For example, *Striga*, *Alectra* and *Orobanchae* infect a wide range of plants in the dry tropics and subtropics. In Africa, five of the most economically important *Striga* species, such as *Striga hermonthica*, *Striga asiatica*, *Striga forbesii*, *Striga aspera* and *Striga gesnerioides*, affect the production of sorghum, millet, maize, sugarcane and cowpea resulting in losses of up to USD 1 billion annually affecting >100 million farmers [4, 11]. Moreover, hundred thousand hectares in Europe, Russia, Middle East, China, Cuba and parts of the USA are infested with *Orobanchae* species [13].

Efforts to control destructive parasitic plants such as *Striga* and *Orobanchae* have been hampered by the complicated association between the parasites and their hosts. Several control methods, including crop rotation, hand-pulling, biological control using *Fusarium oxysporum* [14] and chemicals such as ethylene gas [15], have been tried, but their effectiveness is limited by the size of infested areas, inefficient delivery systems and costs. Therefore, development of crops resistant against these hazardous parasitic plants is proposed to be the only viable strategy [2]. Understanding the molecular mechanism allowing parasitic plants to parasitize host plants is an important step toward generating new resistant crops. Combining advanced technologies such as next-generation sequencing and bioinformatics has allowed us to perform genome-wide gene expression analysis (transcriptomics) on non-model organisms to investigate the molecular mechanisms of host–parasite interactions [16]. In this review, we highlight recent key findings on parasitic plants obtained from transcriptomics studies and the unique aspects that allow parasitic plants to support their heterotrophic life cycle.

## Haustorium

### What is the haustorium of parasitic plant?

The haustorium is a key multicellular organ shared across all parasitic plants, providing physical and physiological bridges between hosts and parasites (Figure 1A) [17]. Root parasites such as *Orobanchaceae* species have a terminal/primary haustorium, which develops on the tip of a primary root soon after germination, and also a lateral/secondary haustorium, which forms laterally on the growing roots without loss of root tip growth. These two types of haustoria show similar anatomic structures and physiological roles. During haustorial development, cells in the tip or lateral side of the root start expansion and division making a globular-shaped organ [18–20]. Concomitantly, the epidermal cells differentiate to hair cells, recognized as haustorial hairs [21]. After penetration into host tissue, host and parasite vessels are connected via the so-called ‘xylem bridge’ (Figure 1B) [22]. A ‘hyaline body’ occupies the central haustorial parts surrounding the xylem cells. This hyaline body has distinctive parenchymatous cells with dense cytoplasm and characteristic extracellular deposits, which may have a function associated with nutrient translocation [23, 24]. Anticlinally elongated epidermal cells, designated as palisade cells, line up at parasite–host interface of the haustorium [25]. *Striga* species also develop a specialized structure called

‘oscula’, which intrusively penetrates into host vessel elements to absorb water and nutrient [26].

By contrast, the stem parasite *Cuscuta* has only rudimentary root system, which disappears a few days after germination and is replaced by haustoria, which develop from the differentiated stem. Development of a haustorium is preceded by a prehaustorium whose initiation is manifested by cell division at sites adjacent to the vascular bundle. In this prehaustorium stage, the vascular proximal cells are dividing, while the tip of the prehaustoria has a zone of large elongated cells, eventually differentiating into searching hyphae [27]. The searching hyphae transforms into xylem or phloem structures depending on the cell types that hyphae encounter on host penetration [28]. Thus, similar but distinct structures underpin the parasitic functions of haustoria in root and stem parasites.


Most facultative parasites are generalists in host specificity, but some obligate parasites only parasitize a limited range of host species [29]. For instance, *S. asiatica* and *S. hermonthica* parasitize only monocots in natural conditions. Nevertheless, they can penetrate but not completely parasitize eudicots, whereas *S. gesnerioides* parasitize eudicots, especially cowpea. Given the different types of incompatibilities shown in interactions between *S. hermonthica* and nonhost species, it is likely that host range is determined by multiple mechanisms [9]. Interestingly, *S. hermonthica* rarely forms haustoria when germinated alongside *Phtheirospermum japonicum*, a facultative parasite in *Orobanchaceae*, suggesting the existence of mechanisms for recognizing closely related parasitic species [9].

### Auxins functions in haustorium

Auxins are crucial in the development of almost all organs of plants [30]. Exogenous application of auxin positively affects haustorium numbers, while disturbing auxin flux using auxin efflux or activity inhibitors results in reduction of haustorium number in *Triphysaria versicolor* and in *Phelipanche aegyptiaca* [31, 32]. Series of root tip dissection experiments have demonstrated that local auxin accumulation is involved in haustorium initiation in *T. versicolor* [32]. Expression of auxin responsive IAA2 promoter in hairy root of *T. versicolor* is further evidence of this local auxin accumulation [32]. A transcriptome study in the stem parasite *Cuscuta pentagona* revealed that genes associated with polar auxin transport activity are enriched in the haustoria stage as compared with stems or seedlings [33]. These data suggest that auxin maxima are established to induce haustoria formation in both root and stem parasites. In addition to auxin, an ethylene response is detected in *T. versicolor* [32] and gibberellic acid- and strigolactone-related genes are upregulated in the haustorium of *C. pentagona* [33]. These observations imply cross talks among multiple phytohormones in haustorial development.

### Haustorial genes

Currently, genes functioning in haustoria of parasitic plants are being discovered by use of molecular genetics tools in combination with transcriptomics. For example, transcriptomes combined with laser capture micro-dissection approaches successfully identified gene expression in haustorial cells in *T. versicolor* [34]. *Triphysaria versicolor* is a generalist parasite with a wide host range [29]. Comparison of gene expression in the *T. versicolor* haustoria penetrating different hosts, such as the monocot *Zea mays* and the eudicot *Medicago truncatula*, identified gene sets that are expressed commonly and differently between



**Figure 1.** Haustorium of parasitic plant. (A) *Striga hermonthica* parasitizing rice roots. (B) Cross section of *S. hermonthica* haustorium penetrating rice root. The section was stained with Safranin O and Fast Green. Px, parasite xylem; Hx, host xylem; HB, hyaline body; COR, cortical parenchyma; INT, interface between the haustorium and the host root. Bar is 500  $\mu$ m.

host species. Although  $\sim 30\%$  of genes highly expressed at the host-parasite interface are unknown, *Expansin* genes are expressed in haustorial cells in a host-specific manner [34]. *Expansins* act nonenzymatically to loosen cell walls;  $\beta$ -*expansins* have an activity specific to grass species, whereas  $\alpha$ -*expansins* target eudicots and other monocots [35–37]. Consistent with their protein functions, expression of  $\beta$ -*expansin* is higher in the *T. versicolor*-infecting *Z. mays* as compared with that in *T. versicolor*-infecting *M. truncatula* and  $\alpha$ -*expansin* expression levels are similar in both interactions. These results suggest that the generalist parasites may leverage distinct genes to parasitize different host plants.


Alakonya et al. [38] found that the *C. pentagona* SHOOT MERISTEMLESS-like (*CpSTM*) gene is highly expressed at prehaustorial stage in *C. pentagona* with a tobacco host. Silencing signals of small interfering RNA (siRNA) can be transmitted through parasite-host attachment as shown in root parasites [39, 40]. Using the cross-species movement of siRNA revealed that knocking down of *CpSTM* disrupts dodder growth by inducing defects in haustorium connection, development and establishment as well as misdirecting growth of searching hyphae [38]. As *STM* is known to regulate shoot development in other plant species [41], this finding contrasts previous hypothesis that dodder haustorium has root origin. Thus, involvement of *CpSTM* in haustorium development programs suggests that the haustorium has a mixed nature of shoot and root origin by co-opting both developmental programs.

## Plant-plant interaction

### Recognition of host

It has been known for many years that Orobanchaceae plants generally develop haustoria only when grown in the presence of other plants (Figure 2A) [2]. The first chemicals identified as haustorium-inducing factors were the flavonoids xenogonin A

and xenogonin B [42], while the first haustorium-inducing factor isolated from actual host roots is 2,6-dimethoxy-p-benzoquinone (DMBQ) [43]. Biochemical analysis of DMBQ and its analogs suggests that semiquinone intermediates, formed during redox cycling between quinone and hydroquinone states, initiate haustorium development [44]. Quinone redox changes are catalyzed by quinone oxidoreductases. Quinone oxidoreductases *TvQR1* and *TvQR2*, which were isolated from the facultative parasite *T. versicolor*, are upregulated after treatment with DMBQ and other quinones [45]. RNA interference experiments show that *TvQR1* is necessary for haustorium development [46]. Given that the *TvQR1* enzyme is in the  $\zeta$ -crystallin family of quinone oxidoreductases [46], the haustorium signaling system could be redox-activated by radicals produced by the *TvQR1* reaction with haustorium-inducing factors. Transcriptomics on DMBQ treatment identified genes transcriptionally regulated by haustorium-inducing factors including *TvQR1/2* [47]. Among the identified genes, *TvPirin*, which is an ortholog associated with various environmental signaling [48, 49], was shown to be necessary for haustorium development in *T. versicolor* [50]. Furthermore, a population genetic study revealed that *TvQR1* exhibited remarkably higher molecular diversity and more recombination events than *TvPirin*, suggesting that *TvQR1* and *TvPirin* might have evolved highly distinct roles for haustorium formation [51]. Taken together, these findings point to the current model for haustorium initiation where parasitic plants trigger the host tissue to produce DMBQ, and the derived DMBQ then induces a parasite quinone oxidoreductase, which converts DMBQ into the active single-electron free radical form with the suitable redox potential for haustorium induction (Figure 2A). Notably it is known that DMBQ can induce haustorial formation in *Triphysaria* species, *Striga* species, *Agalinis purpurea* and *P. japonicum* [19, 43–45, 52, 53]. More detailed stage-specific transcriptomics will allow us to unveil the gene regulatory network that operates during haustorium formation under DMBQ activation.



**Figure 2.** Parasite–host interactions. (A) Model for root parasite–host interactions to complete parasitizing. (B) Model for macromolecule trafficking between parasite and host cells. Because data were obtained from different root parasites and host plants, plant names are indicated in parentheses for each gene/pathway/transcript.

### Exchange of molecules


Because parasitic plants directly connect to host plants through a continuous vascular system (and plasmodesmata in some species like *Striga* and *Cuscuta* [54–56]), movement of molecules between these plants has been reported (Figure 2B) [7]. Currently, there is evidence that parasitic plants acquire various types of macromolecules, such as DNA [5, 57–64], mRNA [65, 66], siRNA [38–40], protein [54, 67, 68] and also viruses [54, 69, 70], from their host plants. Interestingly, transcriptome analysis using *Cuscuta* and its hosts *Arabidopsis* and tomato revealed ‘genomic-scale’ exchanges of mRNA between parasitic and host plants [71]. The uptake and distribution of *Arabidopsis* mRNA into *Cuscuta* varies depending on the mRNA under investigation, indicating multiple routes of RNA trafficking or selective mechanism for mobile mRNA [72]. These findings raise the possibility that the movement of non-nutrient solutes may function physiologically and ecologically in parasite–host interactions [73]. For example, movement of solutes may create osmotic pressure to take up water. In addition, as siRNA can be moved between parasite and host plants [39, 40], it is possible that exchange of small RNA is involved in regulation of gene expression in both parasite and host at the transcriptional level by DNA methylation, as well as at the posttranscriptional level by direct mRNA interference [74]. A combination of detailed transcriptomics, bioinformatics and functional analyses could be a powerful tool to assess whether parasitic plants have evolved mechanisms to exchange molecules to their advantage.

In the course of evolution, nucleic acids were not only transferred from host plants but also inserted into the parasite genomes (Figure 2B). This is termed as horizontal gene transfer (HGT) and is defined as the movement of genetic material between species other than by descent. HGT of both mitochondrial and nuclear genes are reported in parasitic plants [2, 57], although HGT of mitochondrial genes might be much higher than that of nuclear genes [58]. Interestingly, a nuclear gene in *Striga*, which was derived from grass via HGT, was discovered to lack introns and has the remnants of a poly-A sequence, suggesting that the transfer occurred through an mRNA intermediate [57]. Transcriptome data capturing the dynamics of mRNA can be used for phylogenomic approaches to identify HGT at the system level. Using such an approach, Xi et al. [59] examined transcriptome data of both the parasite (*Rafflesia*)

and its host (*Tetrastigma*) to reveal the unidirectional host-to-parasite gene transfers. *Rafflesia* HGTs represent a wide range of cellular functions including respiration, metabolism and protein turnover, and were expressed at levels comparable with vertical gene transfer (VGT) [59]. This indicates that HGT in parasitic plants is not just a by-product of molecule movement through vascular connection but might be mechanically and biologically meaningful. Notably, VGT in *Rafflesia* shows host-like codon usage properties compared with their closest relatives [59]. This could be interpreted as suggesting that VGT may have evolved convergently to match the translational requirements of their host, thus making a suitable cellular environment for promoting and/or maintaining HGT in the parasitic plant [59]. Recent study revealed that *P. aegyptiaca* and related parasitic species and even distantly related parasitic species *C. pentagona* have obtained *albumin 1* KNOTTIN-like genes from legumes through separate HGT events [75]. This suggests that certain genes may have been repeatedly captured by parasitic plants.

### Loss of photosynthesis

Although hemiparasites still carry out photosynthesis to some extent, holoparasites completely rely on the host producing carbohydrates. In holoparasites, therefore, photosynthesis-related genes are no longer required, and may become pseudogenes and be deleted, resulting in a functional and physical reduction of the plastid genome [5, 10, 76–78]. Despite the fact that ~90% of photosynthesis-related genes are in the nuclear genome [79], the nuclear genome and transcriptome remain largely unexplored to date. Most recent advances in developing high-throughput RNA-seq allow us to readily generate the transcriptome data for multiple tissues and species [80]. These technical advances were used for multiple species in the parasitic plant family Orobanchaceae [81, 82] and also for different stages of the whole life cycle of *C. pentagona* [33]. The Parasitic Plant Genome Project has sequenced transcripts from three parasitic species, including facultative parasite *T. versicolor*, obligate hemiparasite *S. hermonthica* and obligate holoparasite *P. aegyptiaca*, and a nonparasitic relative in the Orobanchaceae [82]. Bioinformatic analysis of transcriptomes generated from the above-ground



**Figure 3.** Loss of photosynthesis. (A) Expression of chlorophyll biosynthesis and photosynthesis-related genes in hemi- and holoparasites. Despite the absence of photosynthesis in holoparasite *Orobanche*, they still express intact genes in chlorophyll biosynthesis pathway. (B) Developmental expression patterns of photosynthesis- and transport-related genes in *Cuscuta haustoria*. The complementary expression pattern indicates that nutrients obtained from the transport activity might stimulate the decrease in expression of genes related to photosynthesis.

tissue samples reveal that the holoparasite *Orobanche aegyptiaca* lost expression of photosynthesis-related genes but surprisingly retains an intact expressed and selectively constrained chlorophyll synthesis pathway [81]. This raises an interesting question regarding the role of chlorophylls in the holoparasite, especially if chlorophylls are being produced but are not participating in photosynthesis (Figure 3A). The chlorophyll intermediates, like protochlorophyllide, may be involved in signaling rather than photosynthesis [81]. Another holoparasite *Cuscuta* also shows overall low expression of photosynthesis-related genes [33]; however, detailed transcriptomics using different developmental stages of *Cuscuta* revealed the temporal regulation of photosynthesis-

related gene expression, where there is increased expression of genes related to transporter activity and reduced expression of genes related to photosynthesis with progression of plant parasitism [33]. This suggests that after successful parasitism, *Cuscuta* acquires its nutrients from the host plant mostly through haustorial transport, and reduces photosynthesis to a minimal level (Figure 3B). These findings are consistent with the characterization of plastid genomes that show gene losses and increased substitution rates in parasitic plant plastid genomes [5, 10, 76–78]. Detailed analysis, for instance, using comparative transcriptomics with close relatives [69, 78, 83], could reveal the process by which the photosynthesis machinery has been lost during evolution.

## Future directions

Technologies of RNA-seq and bioinformatics analyses have revealed the molecular mechanisms underlying the uniqueness of parasitic plants such as evolving a haustorium, modifying the plant–plant interactions and shutdown of photosynthesis. Technical advances in transformation have also facilitated research in parasitic plants. Transient transformation method for facultative parasites *P. japonicum* [19] and *T. versicolor* [84] and virus-induced gene silencing for *S. hermonthica* [85] have been developed and used for functional analysis. Using transgenic host plants to silence genes in parasitic plants is a distinctive approach in the study of parasite–host interaction that can be applied for improving crops [40, 86]. High-throughput transformation technique using hairy root transformation of host plants is also useful to investigate the function of genes in hosts responding to parasitic plants [87]. However, the stable and heritable transformation, which is necessary for performing detailed molecular genetics of parasitic plants, has not been established yet.

There still remain key questions for the current parasitic plant research field: What determines the uniqueness of parasitic plants? As certain genes show unique characteristics in molecular evolution rate [88], do parasitic plants have unique genome properties to evolve their uniqueness? Or as current evolutionary developmental biology suggests [89], do parasitic plants have conventional genome properties but co-opt preexisted toolkit genes to evolve their uniqueness? To answer these questions, genome sequencing of parasitic plants is necessary in addition to the transcriptomics reviewed in this article. Sequencing parasitic plant genomes and comparing them with those of nonparasitic plants should allow us to determine the extent of conservation of gene content, genomic sequences and other genomic information, including ‘functional elements’ such as non-protein-coding loci, transcription start sites, regulatory sequences, chromatin accessibility and histone modification patterns [90–92] in the parasitic plant genomes. In addition, as parasitism has evolved independently >11 times [5], there may be a common molecular mechanism for parasitism in angiosperms. Interestingly, recent studies reveal that independent lineages have leveraged similar molecular pathways in the convergent evolution seen in electric organs of vertebrates and bioluminescent organs of squids [93, 94]. Similar to this case, there might be a few key genetic components that can induce parasitism in the ancestral angiosperm genome. We believe future comparative genomics across phylogenetically distant parasitic plants may reveal the answer to this question. These future functional genomic studies on parasitic plants will provide new concepts for understanding the evolutionary novelty and heterotrophic ability of parasitic plants. By extension, this will allow us to improve agricultural crops as well as solve the agricultural/economic problem of damage by parasitic plants.

### Key points

- Uniqueness of parasitic plants is characterized by evolving a haustorium, which facilitates plant–plant interactions, and reducing own photosynthesis.
- Transcriptomics with cellular resolution identified key genes functioning in the haustorium.
- Transcriptomics demonstrated ‘genomic-scale’ exchanges of mRNA between parasitic and host plants.
- Transcriptomics revealed the process by which the photosynthesis machinery has been lost in the course of evolution.

## Acknowledgements

We apologize to colleagues whose work could not be highlighted because of space limitations. We thank Thomas Spallek, Juliane Ishida, Takanori Wakatake and Songkui Cui for critical reading of the manuscript.

## Funding

This work is partially supported by MEXT KAKENHI (No. 24228008 to K.S., No. 25114521, 25711019 and 25128716 to S.Y.), JSPS Postdoctoral Fellowship (to J.M.M.) and the RIKEN Special Postdoctoral Researchers Program (to Y.I.).

## References

1. Rispaill N, Dita MA, Gonzalez-Verdejo C, et al. Plant resistance to parasitic plants: molecular approaches to an old foe. *New Phytol* 2007;173:703–12.
2. Westwood JH, Yoder JI, Timko MP, et al. The evolution of parasitism in plants. *Trends Plant Sci* 2010;15:227–35.
3. Press MC, Phoenix GK. Impacts of parasitic plants on natural communities. *New Phytol* 2005;166:737–51.
4. Spallek T, Mutuku M, Shirasu K. The genus *Striga*: a witch profile. *Mol Plant Pathol* 2013;14:861–9.
5. Barkman TJ, McNeal JR, Lim SH, et al. Mitochondrial DNA suggests at least 11 origins of parasitism in angiosperms and reveals genomic chimerism in parasitic plants. *BMC Evol Biol* 2007;7:248.
6. Yoshida S, Shirasu K. Plants that attack plants: molecular elucidation of plant parasitism. *Curr Opin Plant Biol* 2012;15:708–13.
7. Leblanc M, Kim G, Westwood JH. RNA trafficking in parasitic plant systems. *Front Plant Sci* 2012;3:203.
8. Yoder JI, Scholes JD. Host plant resistance to parasitic weeds; recent progress and bottlenecks. *Curr Opin Plant Biol* 2010;13:478–84.
9. Yoshida S, Shirasu K. Multiple layers of incompatibility to the parasitic witchweed, *Striga hermonthica*. *New Phytol* 2009;183:180–9.
10. dePamphilis CW, Palmer JD. Loss of photosynthetic and chlororespiratory genes from the plastid genome of a parasitic flowering plant. *Nature* 1990;348:337–9.
11. Pennisi E. Armed and dangerous. *Science* 2010;327:804–5.
12. Ezeaku IE, Gupta SC. Development of sorghum populations for resistance to *Striga hermonthica* in the Nigerian Sudan Savanna. *Afr J Biotechnol* 2004;3:324–9.
13. Musselman L, Press M. *Introduction to Parasitic Plants*. London: Chapman and Hall, 1995.
14. Yonli D, Hess DE, Abbasher AA, et al. Biological control of witch weed in fields of Burkina Faso using isolates of *Fusarium oxysporum*. *Afr Crop Sci J* 2005;13:41–7.
15. Inverson RD, Westbrooks RG, Eplee RE, et al. *Overview and Status of the Witchweed (Striga asiatica) Eradication Program in the Carolinas*. Washington: American Chemical Society, 2011.
16. Wang Z, Gerstein M, Snyder M. RNA-Seq: a revolutionary tool for transcriptomics. *Nat Rev Genet* 2009;10:57–63.
17. Kujit J. *The Biology of Parasitic Flowering Plants*. Berkeley: University of California Press, 1969.
18. Baird WV, Riopel JL. Experimental studies of haustorium initiation and early development in *Agalinis purpurea* (L.) Raf. (*Scrophulariaceae*). *Am J Bot* 1984;71:803–14.

19. Ishida JK, Yoshida S, Ito M, et al. *Agrobacterium rhizogenes*-mediated transformation of the parasitic plant *Phtheirospermum japonicum*. *PLoS One* 2011;**6**:e25802.
20. Riopel JL, Musselman L. Experimental initiation of haustoria in *Agalinis purpurea*. *Am J Bot* 1979;**66**:570–5.
21. Baird WV, Riopel JL. Surface characteristics of root haustorial hairs of parasitic *Scrophulariaceae*. *Bot Gaz* 1985;**146**:63–9.
22. Heide-Jorgensen HS, Kuijt J. The haustorium of the root parasite *Triphysaria* (*Scrophulariaceae*), with special reference to xylem bridge ultrastructure. *Am J Bot* 1995;**82**:782–97.
23. Pielach A, Leroux O, Domozych DS, et al. Arabinogalactan protein-rich cell walls, paramural deposits and ergastic globules define the hyaline bodies of rhinanthoid *Orobanchaceae* haustoria. *Ann Bot* 2014;**114**:1359–73.
24. Visser JH, Dorr I, Kollmann R. The hyaline body of the root parasite *Alectra-orobanchoides* benth (*scrophulariaceae*) - its anatomy, ultrastructure and histochemistry. *Protoplasma* 1984;**121**:146–56.
25. Heide-Jorgensen HS, Kuijt J. Epidermal derivatives as xylem elements and transfer cells: a study of the host-parasite interface in two species of *Triphysaria* (*Scrophulariaceae*). *Protoplasma* 1993;**174**:173–83.
26. Dorr I. How *Striga* parasitizes its host: a TEM and SEM study. *Ann Bot* 1997;**79**:463–72.
27. Lee KB. Structure and development of the upper haustorium in the parasitic flowering plant *Cuscuta japonica* (*Convolvulaceae*). *Am J Bot* 2007;**94**:737–45.
28. Vaughn KC. Conversion of the searching hyphae of dodder into xylem and phloem hyphae: a cytochemical and immunocytochemical investigation. *Int J Plant Sci* 2006;**167**:1099–114.
29. Musselman L. The biology of *Striga*, *Orobanche*, and other root-parasitic weeds. *Annu Rev Phytopathol* 1980;**18**:463–89.
30. Finet C, Jaillais Y. Auxology: when auxin meets plant evolution. *Dev Biol* 2012;**369**:19–31.
31. Bar-Nun N, Sachs T, Mayer AM. A role for IAA in the infection of *Arabidopsis thaliana* by *Orobanche aegyptiaca*. *Ann Bot* 2008;**101**:261–5.
32. Tomilov AA, Tomilova NB, Abdallah I, et al. Localized hormone fluxes and early haustorium development in the hemiparasitic plant *Triphysaria versicolor*. *Plant Physiol* 2005;**138**:1469–80.
33. Ranjan A, Ichihashi Y, Farhi M, et al. *De novo* assembly and characterization of the transcriptome of the parasitic weed *Cuscuta pentagona* identifies genes associated with plant parasitism. *Plant Physiol* 2014;**166**:1186–99.
34. Honaas LA, Wafula EK, Yang Z, et al. Functional genomics of a generalist parasitic plant: laser microdissection of host-parasite interface reveals host-specific patterns of parasite gene expression. *BMC Plant Biol* 2013;**13**:9.
35. Cosgrove DJ. Loosening of plant cell walls by expansins. *Nature* 2000;**407**:321–6.
36. Li Y, Jones L, McQueen-Mason S. Expansins and cell growth. *Curr Opin Plant Biol* 2003;**6**:603–10.
37. Sampedro J, Cosgrove DJ. The expansin superfamily. *Genome Biol* 2005;**6**:242.
38. Alakonya A, Kumar R, Koenig D, et al. Interspecific RNA interference of SHOOT MERISTEMLESS-like disrupts *Cuscuta pentagona* plant parasitism. *Plant Cell* 2012;**24**:3153–66.
39. Aly R, Cholakh H, Joel DM, et al. Gene silencing of mannose 6-phosphate reductase in the parasitic weed *Orobanche aegyptiaca* through the production of homologous dsRNA sequences in the host plant. *Plant Biotechnol J* 2009;**7**:487–98.
40. Tomilov AA, Tomilova NB, Wroblewski T, et al. Trans-specific gene silencing between host and parasitic plants. *Plant J* 2008;**56**:389–97.
41. Hay A, Tsiantis M. KNOX genes: versatile regulators of plant development and diversity. *Development* 2010;**137**:3153–65.
42. Steffens JC, Lynn DG, Kamat VS, et al. Molecular specificity of haustorial induction in *Agalinis purpurea* (L.) Raf. (*Scrophulariaceae*). *Ann Bot* 1982;**50**:1–7.
43. Chang M, Lynn DG. The haustorium and the chemistry of host recognition in parasitic angiosperms. *J Chem Ecol* 1986;**12**:561–79.
44. Smith CE, Ruttledge T, Zeng Z, et al. A mechanism for inducing plant development: the genesis of a specific inhibitor. *Proc Natl Acad Sci USA* 1996;**93**:6986–91.
45. Matvienko M, Wojtowicz A, Wrobel R, et al. Quinone oxidoreductase message levels are differentially regulated in parasitic and non-parasitic plants exposed to allelopathic quinines. *Plant J* 2001;**25**:375–87.
46. Bandaranayake PC, Filappova T, Tomilov A, et al. A single-electron reducing quinone oxidoreductase is necessary to induce haustorium development in the root parasitic plant *Triphysaria*. *Plant Cell* 2010;**22**:1404–19.
47. Matvienko M, Torres MJ, Yoder JI. Transcriptional responses in the hemiparasitic plant *Triphysaria versicolor* to host plant signals. *Plant Physiol* 2001;**127**:272–82.
48. Lapić YR, Kaufman LS. The *Arabidopsis* cupin domain protein AtPirin1 interacts with the G protein alpha-subunit GPA1 and regulates seed germination and early seedling development. *Plant Cell* 2003;**15**:1578–90.
49. Warpeha KM, Upadhyay S, Yeh J, et al. The GCR1, GPA1, PRN1, NF-Y signal chain mediates both blue light and abscisic acid responses in *Arabidopsis*. *Plant Physiol* 2007;**143**:1590–600.
50. Bandaranayake PC, Tomilov A, Tomilova NB, et al. The TvPirin gene is necessary for haustorium development in the parasitic plant *Triphysaria versicolor*. *Plant Physiol* 2012;**158**:1046–53.
51. Ngo QA, Albrecht H, Tsuchimatsu T, et al. The differentially regulated genes TvQR1 and TvPirin of the parasitic plant *Triphysaria* exhibit distinctive natural allelic diversity. *BMC Plant Biol* 2013;**13**:28.
52. Jamison DS, Yoder JI. Heritable variation in quinone-induced haustorium development in the parasitic plant *Triphysaria*. *Plant Physiol* 2001;**125**:1870–9.
53. Stranger A, Murphy A, Corbett JM, et al. Changes in patterns of protein-synthesis during haustorial development of *Striga hermonthica* (Del) Benth seedlings. *J Exp Bot* 1995;**46**:277–83.
54. Birschwilks M, Haupt S, Hofius D, et al. Transfer of phloem-mobile substances from the host plants to the holoparasite *Cuscuta* sp. *J Exp Bot* 2006;**57**:911–21.
55. Vaughn KC. Dodder hyphae invade the host: a structural and immunocytochemical characterization. *Protoplasma* 2003;**220**:189–200.
56. Dorr I. *Sieve elements in haustoria of parasitic angiosperms*. Berlin, Germany: Springer-Verlag, 1990.
57. Yoshida S, Maruyama S, Nozaki H, et al. Horizontal gene transfer by the parasitic plant *Striga hermonthica*. *Science* 2010;**328**:1128.
58. Xi Z, Wang Y, Bradley RK, et al. Massive mitochondrial gene transfer in a parasitic flowering plant clade. *PLoS Genet* 2013;**9**:e1003265.
59. Xi Z, Bradley RK, Wurdack KJ, et al. Horizontal transfer of expressed genes in a parasitic flowering plant. *BMC Genomics* 2012;**13**:227.

60. Park JM, Manen JF, Schneeweiss GM. Horizontal gene transfer of a plastid gene in the non-photosynthetic flowering plants *Orobanche* and *Phelipanche* (Orobanchaceae). *Mol Phylogenet Evol* 2007;**43**:974–85.
61. Mower JP, Stefanovic S, Young GJ, et al. Plant genetics: gene transfer from parasitic to host plants. *Nature* 2004;**432**:165–6.
62. Mower JP, Stefanovic S, Hao W, et al. Horizontal acquisition of multiple mitochondrial genes from a parasitic plant followed by gene conversion with host mitochondrial genes. *BMC Biol* 2010;**8**:150.
63. Davis CC, Wurdack KJ. Host-to-parasite gene transfer in flowering plants: phylogenetic evidence from Malpighiales. *Science* 2004;**305**:676–8.
64. Davis CC, Anderson WR, Wurdack KJ. Gene transfer from a parasitic flowering plant to a fern. *Proc Biol Sci* 2005;**272**:2237–42.
65. David-Schwartz R, Runo S, Townsley B, et al. Long-distance transport of mRNA via parenchyma cells and phloem across the host-parasite junction in *Cuscuta*. *New Phytol* 2008;**179**:1133–41.
66. Roney JK, Khatibi PA, Westwood JH. Cross-species translocation of mRNA from host plants into the parasitic plant dodder. *Plant Physiol* 2007;**143**:1037–43.
67. Haupt S, Oparka KJ, Sauer N, et al. Macromolecular trafficking between *Nicotiana tabacum* and the holoparasite *Cuscuta reflexa*. *J Exp Bot* 2001;**52**:173–7.
68. Aly R, Hamamouch N, Abu-Nassar J, et al. Movement of protein and macromolecules between host plants and the parasitic weed *Phelipanche aegyptiaca* Pers. *Plant Cell Rep* 2011;**30**:2233–41.
69. Jiang L, Wijeratne AJ, Wijeratne S, et al. Profiling mRNAs of two *Cuscuta* species reveals possible candidate transcripts shared by parasitic plants. *PLoS One* 2013;**8**:e81389.
70. Gal-On A, Naglis A, Leibman D, et al. Broomrape can acquire viruses from its hosts. *Phytopathology* 2009;**99**:1321–9.
71. Kim G, LeBlanc ML, Wafula EK, et al. Genomic-scale exchange of mRNA between a parasitic plant and its hosts. *Science* 2014;**345**:808–11.
72. LeBlanc M, Kim G, Patel B, et al. Quantification of tomato and Arabidopsis mobile RNAs trafficking into the parasitic plant *Cuscuta pentagona*. *New Phytol* 2013;**200**:1225–33.
73. Smith JD, Mescher MC, De Moraes CM. Implications of bioactive solute transfer from hosts to parasitic plants. *Curr Opin Plant Biol* 2013;**16**:464–72.
74. Pumphlin N, Voinnet O. RNA silencing suppression by plant pathogens: defence, counter-defence and counter-counter-defence. *Nat Rev Microbiol* 2013;**11**:745–60.
75. Zhang Y, Fernandez-Aparicio M, Wafula EK, et al. Evolution of a horizontally acquired legume gene, albumin 1, in the parasitic plant *Phelipanche aegyptiaca* and related species. *BMC Evol Biol* 2013;**13**:48.
76. Funk HT, Berg S, Krupinska K, et al. Complete DNA sequences of the plastid genomes of two parasitic flowering plant species, *Cuscuta reflexa* and *Cuscuta gronovii*. *BMC Plant Biol* 2007;**7**:45.
77. McNeal JR, Arumugunathan K, Kuehl JV, et al. Systematics and plastid genome evolution of the cryptically photosynthetic parasitic plant genus *Cuscuta* (Convolvulaceae). *BMC Biol* 2007;**5**:55.
78. Wicke S, Muller KF, de Pamphilis CW, et al. Mechanisms of functional and physical genome reduction in photosynthetic and nonphotosynthetic parasitic plants of the broomrape family. *Plant Cell* 2013;**25**:3711–25.
79. Martin W, Herrmann RG. Gene transfer from organelles to the nucleus: how much, what happens, and Why? *Plant Physiol* 1998;**118**:9–17.
80. Kumar R, Ichihashi Y, Kimura S, et al. A high-throughput method for Illumina RNA-seq library preparation. *Front Plant Sci* 2012;**3**:202.
81. Wickett NJ, Honaas LA, Wafula EK, et al. Transcriptomes of the parasitic plant family Orobanchaceae reveal surprising conservation of chlorophyll synthesis. *Curr Biol* 2011;**21**:2098–104.
82. Westwood J, dePamphilis CW, Das M, et al. The parasitic plant genome project: new tools for understanding the biology of *Orobanche* and *Striga*. *Weed Sci* 2012;**60**:295–306.
83. Braukmann T, Kuzmina M, Stefanovic S. Plastid genome evolution across the genus *Cuscuta* (Convolvulaceae): two clades within subgenus *Grammica* exhibit extensive gene loss. *J Exp Bot* 2013;**64**:977–89.
84. Tomilov A, Tomilova N, Yoder JI. *Agrobacterium tumefaciens* and *Agrobacterium rhizogenes* transformed roots of the parasitic plant *Triphysaria versicolor* retain parasitic competence. *Planta* 2007;**225**:1059–71.
85. Kirigia D, Runo S, Alakonya A. A virus-induced gene silencing (VIGS) system for functional genomics in the parasitic plant *Striga hermonthica*. *Plant Methods* 2014;**10**:16.
86. Bandaranayake PC, Yoder JI. Trans-specific gene silencing of acetyl-CoA carboxylase in a root-parasitic plant. *Mol Plant Microbe Interact* 2013;**26**:575–84.
87. Runo S, Macharia S, Alakonya A, et al. *Striga* parasitizes transgenic hairy roots of *Zea mays* and provides a tool for studying plant-plant interactions. *Plant Methods* 2012;**8**:20.
88. Bromham L, Cowman PF, Lanfear R. Parasitic plants have increased rates of molecular evolution across all three genomes. *BMC Evol Biol* 2013;**13**:126.
89. Peter IS, Davidson EH. Evolution of gene regulatory networks controlling body plan development. *Cell* 2011;**144**:970–85.
90. Consortium EP, Birney E, Stamatoyannopoulos JA, et al. Identification and analysis of functional elements in 1% of the human genome by the ENCODE pilot project. *Nature* 2007;**447**:799–816.
91. Gerstein MB, Lu ZJ, Van Nostrand EL, et al. Integrative analysis of the *Caenorhabditis elegans* genome by the modENCODE project. *Science* 2010;**330**:1775–87.
92. mod EC, Roy S, Ernst J, et al. Identification of functional elements and regulatory circuits by *Drosophila* modENCODE. *Science* 2010;**330**:1787–97.
93. Gallant JR, Traeger LL, Volkening JD, et al. Nonhuman genetics. Genomic basis for the convergent evolution of electric organs. *Science* 2014;**344**:1522–5.
94. Pankey MS, Minin VN, Imholte GC, et al. Predictable transcriptome evolution in the convergent and complex bioluminescent organs of squid. *Proc Natl Acad Sci USA* 2014;**111**:E4736–42.